MOTRIN® Professional Dosing Information for Infants & Children

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving MOTRIN*. **A healthcare professional should be consulted for dosing in children under the age of six months.**



| | Your chil | Your child's weight and age | | | | | | | |
|---|------------------------------|--|------------------------|----------------------|-----------------------------------|----------------------|---------------------------|---------------------|--|
| Always ask your healthcare professional which product is right for your child. | Age 0-5 mos //eight 6-11 lbs | 6-11 mos 12-17 lbs | 12-23 mos 18-23 lbs | 2-3 yrs 24-35 lbs | 4-5 yrs 36-47 lbs | 6-8 yrs 48-59 lbs | 9-10 yrs 60-71 lbs | 11 yrs 72-95 lbs | |
| Concentrated MOTRIN® Infants' Drops Berry & Berry dye-free flavors Active Ingredient: Ibuprofen 50mg (in each 1.25mL) Use only as directed. | - | 1.25mL | 1.875mL | - | - | - | - | - | |
| Children's MOTRIN® Suspension Berry, Grape, Tropical Punch, Bubblegum & Berry dye-free flavors Active Ingredient: Ibuprofen 100mg (in each 5mL or 1 tsp) Use only as directed. | on – | _ | _ | 1 tsp or 5mL | 1 ¹ /2 tsp or 7.5mL | 2 tsp or 10mL | 2 ½ tsp or 12.5mL | 3 tsp or 15mL | |
| MOTRIN® Junior Strength Easy-to-Swallow Caplets Active Ingredient: Ibuprofen 100mg (in each caplet) Use only as directed. | - | _ | - | - | - | 2 caplets | 2 ¹ /2 caplets | 3 caplets | |
| MOTRIN® Junior Strength Chewable Caplets Orange & Grape flavors Active Ingredient: Ibuprofen 100mg (in each tablet) | - | - | - | 1 tablet | 11/2 tablets tsp = teaspoon* | 2 tablets | 2 ¹ /2 tablets | 3 tablets | |
| ı | | Take every 6-8 hours as needed. Do not exceed more than 4 doses in 24 hours. | | | | | | | |

Your child's woight and ago

Important Instructions for Proper Use

- Read and follow the label instructions on all MOTRIN® products
- Take every 6-8 hours as needed
- Do not exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID (prescription or nonprescription)
- · Keep all medicines out of the reach of children
- *Use only the dosing device that comes with a specific product
 - Household items such as measuring spoons are less accurate
- Do NOT administer adult medicines to children
- Concentrated MOTRIN® Infants' Drops are more concentrated than Children's MOTRIN® Liquids. The Concentrated Infants' Drops have been specifically designed for use with enclosed dosing device.
- Children's MOTRIN* Liquids are less concentrated than Concentrated MOTRIN* Infants' Drops. The Children's MOTRIN* Liquids have been specifically designed for use with the enclosed measuring cup.
- The OTC dosing: the recommended dose is 7.5mg/kg every 6-8 hours. The recommended daily dose is 30mg/kg.
- If you have any questions, contact your healthcare professional or call 1-877-895-3665

- MOTRIN® contains ibuprofen

©McNEIL-PPC Inc. 2009 CTCM-0058